

PROCEEDINGS
OF THE
BIOLOGICAL SOCIETY OF WASHINGTON

PUPA OF *NEOMOCHTHERUS ANGUSTIPENNIS*
(HINE), WITH NOTES ON FEEDING HABITS OF
ROBBER FLIES AND A REVIEW OF PUBLICATIONS
ON MORPHOLOGY OF IMMATURE STAGES
(DIPTERA: ASILIDAE)

By L. V. KNUTSON


*Systematic Entomology Laboratory,
Agricultural Research Service, USDA¹*

FEEDING HABITS

Robber flies are one of the most taxonomically diverse and numerically abundant groups of Diptera. As far as known, the adults and some (possibly all) larvae are predators of other insects, especially phytophagous forms. Some adult asilids attack beneficial insects such as honey bees and other Hymenoptera, and some prey on harmful species, such as locusts, grasshoppers, and chafer-beetles. Although these common flies appear to be significant elements of the ecosystem and of practical importance to agriculture, surprisingly little is known about the behavior of the adults. Even less is known about the immature stages.

In addition to many isolated, single-specimen observations of prey, extensive lists of prey of adult Asilidae have been presented (among others) by Adamovic (1964) and Hobby (1930, 1933) in Europe, Bromley (1930) and Fattig (1945) in North America, Carrera and Vulcano (1961) in South America, Cuthbertson (1937) in Africa, and Iwata and Nagatomi (1962) in Japan. No adult asilids are known to be monophagous.

¹ Mail address: c/o National Museum of Natural History, Smithsonian Institution, Washington, D.C. 20560.



gous or even oligophagous. Whereas some robber flies are opportunistic predators and attack most of the abundant, slow-flying, appropriately sized and conspicuous species available (Powell and Stage, 1962), others have more limited prey preferences (Linsley, 1960; Lavigne and Holland, 1969). Apparently many species have hunting zones restricted to certain parts of the habitat (Adamovic, 1963; Lehr, 1969; Zinov'eva, 1959). As is true of most other groups of entomophagous insects, there is little quantitative data on the impact of adult robber flies on populations of their prey.

The most extensive study of the biology of Asilidae (Melin, 1923) includes much information on various aspects of all stages, especially the adults, but little specific data on the critical issues of the food and feeding behavior of the larvae. From the fragmentary data of many earlier authors (but not including the important information of Felt, 1918, and Davis, 1919) and from his own mainly circumstantial evidence, Melin concluded that asilid larvae are principally phytophagous. This opinion, made without knowledge of the complete life cycle of any species, has been repeated in many subsequent general discussions of the biology of Asilidae.

Since Melin's paper, conclusive studies of the feeding habits of larvae belonging to the genera *Hyperechia*, *Mallophora*, and *Promachus* have appeared. Adults of *Hyperechia* are well known as striking mimics of *Xylocopa* bees in the Old World Tropics, and at least the mature larvae of several species of *Hyperechia* are predators of the larvae and pupae of the xylocopid models (Poulton, 1924; Tsacas, et al., 1970). The complete development of the larvae of only three of the 5,000 known species of Asilidae (*Promachus yesonicus* Bigot, *Mallophora media* Clements and Bennett, and *M. ruficauda* (Wiedemann)) has been studied. The larvae of *Promachus yesonicus* are free-living predators of the larvae of Scarabaeidae, especially of *Anomala* spp., in Japan (Kinoshita, 1940). The larvae of *Mallophora media* were reared from larvae of a scarabaeid (*Barybas insulanus* Moser) in Trinidad by Clements and Bennett (1969). The young larva is an ectoparasite and remains in place with its anterior end buried in the in-

tegument of the grub. During later larval life it may remain with the original host or it may live free in the soil. The young larva of *M. ruficauda* is an ectoparasite of *Archohileurus vortex* (Burmeister) (Scarabaeidae); it remains with the host larva and consumes it as it matures (Copello, 1927, 1942). Less detailed reports of the behavior of the larvae of *M. ruficauda* are found in the papers of Crouzel (1965) and De Gavotto (1964).

Other reports on the food and feeding habits of asilid larvae not included in Melin's 1923 review or published subsequently are summarized below. Most of these also indicate that the larvae of Asilidae are predaceous. Felt (1918) reported that larvae of *Promachus fitchii* Osten Sacken are efficient natural enemies of larvae of *Phyllophaga fusca* (Froelich) (Scarabaeidae) in New York. Davis (1919) observed free-living larvae of *Promachus vertebratus* (Say) feeding on white grubs in Wisconsin and Michigan, and he cited unpublished reports of other asilid larvae feeding on scarabaeid larvae. Irwin-Smith (1923) kept first-instar larvae of an Australian species, *Neoaratus hercules* Wiedemann, alive for more than 25 days but the larvae did not feed on, "... pieces of Scarabaeidae larvae . . ." or on earthworms or scraps of meat. Champlain and Knoll (1923) reared adults from larvae and pupae of *Andrenosoma fulvicauda* (Say) found in pupal cells of *Chrysobothris femorata* (Olivier) (Buprestidae) in *Quercus* sp. in Pennsylvania. Osterberger (1930) reared larvae of *Triorla interrupta* (Macquart) on larvae of *Eutheola rugiceps* (LeConte) (Scarabaeidae) in Louisiana. Osterberger stated that, "The feeding (Pl. 24, fig. 1) was accomplished by the larva attaching itself to the white grub in a tender, unprotected region just back of the head. There it remained until all the body juices were sapped. . . ." Reinhard (1938) was unable to induce newly hatched larvae of *Efferia aestuans* (L.) to feed on small *Phyllophaga* grubs or on ant larvae in Illinois. Ritcher (1940) found over 40 larvae of *Diogmites discolor* Loew attacking pupae or in pupal cells containing shrivelled skins of pupae of five species of *Phyllophaga* in Kentucky. Fattig (1944) stated that, "From hundreds



FIG. 1. Larva of *Mallophora* sp. on larva of *Phyllophaga* sp.

of white grubs brought in with the larva of some insect attached . . . ,” he reared the following in Georgia: *Asilus virginicus* Banks, “*Deromyia winthemi* Wiedemann,” *Diogmites discolor* Loew, *D. ternatus* Loew, *Efferia aestuans* (L.), *E. pogonias* (Wiedemann), “*Erax barbatus* Fabricius,” “*Erax interruptus* (L.),” “*Proctacanthus brevipennis* Wiedemann,” *P. longus* (Wiedemann), *P. rufus* Williston, *Promachus rufipes* (Fabricius), and *Tolmerus notatus* Wiedemann. Stower, et al., (1958) found larvae of *Scylaticus* sp. in egg-pods of *Schistocerca gregaria* (Forskål) in Eritrea. Greathead (1963) recorded larvae of *Scylaticus* sp., *Stenopogon* sp., and *Trichardis* sp. from egg-pods of *S. gregaria* in East Africa. Daniels (1966) found larvae (1–1.5 inches long) of an unidentified species of Asilidae in soil with larvae of *Phyllophaga koehleriana* (Saylor) in Texas: the asilid larvae killed and ate the scarabaeid larvae during laboratory rearings. Ecipenko (1967) collected a larva of *Eutolmus rufibarbis* Meigen, “. . . that was

sucking out a larva of the beetle *Maladera japonica* Motschulsky [Scarabaeidae] through a puncture in the head capsule." During laboratory rearings, Ecipenko saw a larva of *E. rufibarbis* kill and eat a larva of *Machimus* sp. (Asilidae), and he also observed cannibalism in a culture of *E. rufibarbis*. Weinberg (1968) obtained hatching of eggs of *Dysmachus fuscipennis* (Meigen) in Romania. The larvae lived for 11 days in containers of different kinds of soil. Weinberg did not state that the larvae fed, although she did mention that some of the larvae molted to the second instar.

The sites and manner of attachment of young *Mallophora* larvae are very similar to those of the larvae of certain tiphiid wasps that are also ectoparasites of scarabaeid larvae. The presence of a pair of bristles on each thoracic segment and of four pairs of elongate, fine bristles on the last body segment and the absence of lateral abdominal spiracles readily distinguish asilid larvae from tiphiid larvae. Figure 1 shows a larva of *Phyllophaga* sp. (Scarabaeidae) (determined by D. M. Anderson) parasitized by young asilid larvae, possibly of the genus *Mallophora*. Several scarabaeid larvae infested in this manner were sent to the Systematic Entomology Laboratory, USDA, for determination by L. Castenada, Salvadorian Coffee Research Institute, San Salvador. The specimens were collected during September 1968, at El Sunza, Dept. Sonsonate, San Salvador. The asilid larvae closely match the description of *Mallophora media* (Clements and Bennett, 1969) and *M. ruficauda* (Copello, 1927, 1942). Mr. A. N. Clements (personal communication), who has seen the specimens, agrees that they are Asilidae, possibly of the genus *Mallophora*. There are seven specimens of *Mallophora ruficauda* in the National Museum of Natural History, Smithsonian Institution, pinned with pupal skins and labeled as follows: 1♂, So. Amer. Par. Labo., Uruguay, near Montevideo, No. 1013, XII.22, H. L. Parker col., Host, Scarabaeidae. 2♀, as above except 1♀ XII.20, P. A. Berry col. and 1♀ XII.26, P. A. Berry col. 1♀, So. Amer. Paras. Lab., Montevideo, No. 561.9, XII.21, Host, Scarabaeidae, Parker and Silveira. 1♀, as above except No. 5618. 1♀, So. Amer. Paras. Lab., Seriano, Uruguay, No. 6262, XII.17,

in soil, Silveira. 1♀ Canelones, Uruguay, pred. Scarabaeidae, 16.I.1942, 561.11, Lot No. 42-8967, H. L. Parker.

MORPHOLOGY OF IMMATURE STAGES

Only about 2 percent of the approximately 5,000 world species of Asilidae are known in any immature stage. Of some 400 total genera, there is information on the immature stages of only 41. Of the 856 species and 82 genera known from North America, immature stages of only 25 and 12, respectively, have been described. Papers on the morphology and biology of immature stages of Asilidae, published to 1918, were listed by Irwin-Smith (1923). Hennig (1952) summarized descriptive data on the immature stages of 64 species in 30 genera. Information on the morphology of immature stages that has been published since Hennig's and Irwin-Smith's summaries (and earlier papers not included by them) is listed in Table 1.

PUPA OF *NEOMOCHTHERUS ANGUSTIPENNIS* (HINE) AND FEATURES OF THE ADULTS

Neomochtherus Osten-Sacken is a typical example of the paucity of information available on the immature stages of Asilidae. In his revision of the genera of Asilidae of the world, Hull (1962) listed 49 species belonging to this genus. The Old World fauna has been studied intensively recently by Tsacas (1968, 1969), who treated 120 Palearctic and Ethiopian species. Apparently only the first-instar larva and pupa of *N. perplexus* Becker (= *N. hungaricus rossicus* Engel?) (Zinov'eva, 1959) and the egg of *N. kivuensis* Tsacas (Tsacas, 1969) have been described. In the Smithsonian Institution there is a male of the rare *N. angustipennis* (Hine) pinned with a pupal skin, which is described below.

Neomochtherus angustipennis (Hine)

Pupa, Figures 2-5.

Greatest length, 14.0 mm; greatest width of thorax, 3.5 mm; greatest width of abdomen, 3.5 mm, tapering to 1.5 mm at greatest width of last abdominal segment. Subshining straw-yellow, spines and processes glistering reddish brown. Head with a pair of terminal anterior antennal

TABLE 1. Recently published descriptions and figures of the immature stages of Asilidae.

LEPTOGASTRINAE		
<i>Leptogaster cylindrica</i> De Geer	E, IL, P ¹	Kazakhstan, U.S.S.R.
<i>Leptogaster salvia</i> Martin	E	Wyoming, U.S.A.
LAPHRIINAE		
<i>Andrenosoma atra</i> Séguy	P	France
<i>Andrenosom bayardi</i> Séguy	P	France
<i>Andrenosoma fulvicauda</i> (Say)	L	Ohio, U.S.A.
<i>Hypererchia bomboidea</i> (Loew)	L, P	Senegal
<i>Laphria</i> sp.	L	Ohio, U.S.A.
<i>Laphria flava</i> L.	L	England
<i>Laphria gilva</i> L.	E	Germany
<i>Laphria gilva</i> L.	E	South Dakota, U.S.A.
<i>Laphria gilva</i> L.	L	England
<i>Laphria gilva</i> L.	P	France
DASYPOGONINAE		
? <i>Dioctria oelandica</i> (L.)	L	Westmoreland, England
<i>Dioctria rufipes</i> (De Geer)	L	England
<i>Dioctria rufipes</i> (De Geer)	L, P	Cheshire, England
<i>Diognites angustipennis</i> Loew	E	Wyoming, U.S.A.
<i>Diognites basalis</i> (Walker)	E	Indiana, U.S.A.
(as <i>Deromyia umbrina</i> Loew)		
<i>Diognites discolor</i> Loew	L, P	Kentucky, U.S.A.

¹ E = egg, L = Mature larva only, unless otherwise indicated, P = pupa.

TABLE 1 (Continued)

<i>Leptarthrus brevirostris</i> Meigen	P	Sussex, England	Parmenter, 1952
<i>Leptarthrus brevirostris</i> Meigen	E	Surrey, England	Stubbs, 1970
<i>Pritchardia hirtipes</i> (Macquart)	E	Chile	Artigas, 1970
<i>Scylaticus</i> sp.	L, P	East Africa	Greathhead, 1963
<i>Stenopogon coyote</i> Bromley	E	Wyoming, U.S.A.	Lavigne, 1963
ASILINAE			
<i>Apoclea helvipes</i> Loew	E, IL	Kazakhstan, U.S.S.R.	Zinov'eva, 1959
<i>Astochia caspica</i> Hermann	E, IL	Kazakhstan, U.S.S.R.	Zinov'eva, 1959
<i>Dysmachus</i> sp.	L	Germany	Brauns, 1954a
<i>Dysmachus fuscipennis</i> (Meigen)	E, I, II L	Rumania	Weinberg, 1968
? <i>Dysmachus trigonus</i> Meigen	P	Germany	Brauns, 1954b
<i>Echthistus rufinervis</i> Meigen	E, IL, P	Kazakhstan, U.S.S.R.	Zinov'eva, 1959
<i>Eccritosisa rubriventris</i> (Macquart)	P	Chile	Artigas, 1970
<i>Efferia helene</i> (Bromley)	E	Wyoming, U.S.A.	Lavigne, 1964
(as <i>E. bicaudata</i> (Hine))			
" "			
<i>Efferia pallidula</i> (Hine)	E	Wyoming, U.S.A.	Lavigne and Holland, 1969
<i>Efferia staminea</i> (Williston)	E	Wyoming, U.S.A.	Lavigne and Holland, 1969
<i>Eremisca vernalis</i> Zinov'eva	E, L	Kazakhstan, U.S.S.R.	Zinov'eva, 1959
<i>Eutolmus implacidus</i> Loew	E	Kazakhstan, U.S.S.R.	Lehr, 1962
<i>Eutolmus rufibarbus</i> Meigen	L, P	Khabarovsk Region, U.S.S.R.	Eciipenko, 1967

TABLE I. (Continued)

ASILINAE		
<i>Machinus atricapillus</i> Fallén	P	Kazakhstan, U.S.S.R.
<i>Machinus gonatistes</i> Zeller	E, L, P	Kazakhstan, U.S.S.R.
<i>Mallophora media</i> Clements and Bennett	I-IV L, P	Trinidad, West Indies
<i>Mallophora ruficauda</i> (Wiedemann)	E, I-IV L, P	Argentina
<i>Mallophorina frustra</i> Pritchard	E	Arizona, U.S.A.
<i>Mallophorina pulchra</i> Pritchard	E	Arizona, U.S.A.
<i>Promachus bastardii</i> (Macquart)	L	Michigan, U.S.A.
<i>Promachus dimidiatus</i> (Curran)	E	Wyoming, U.S.A.
<i>Promachus leontochlaenus</i> Loew	E	Kazakhstan, U.S.S.R.
<i>Promachus vertebratus</i> (Say)	L, P	U.S.A.
<i>Promachus</i> sp., prob. <i>vertebratus</i> (Say)	L	Ohio, U.S.A.
<i>Neomochtherus kivuensis</i> Tsacas	E	Lake Kivu, Congo
<i>Neonoclutherus perplexus</i> Becker (= <i>N. hungaricus rossicus</i> Engel ?)	E, IL, P	Kazakhstan, U.S.S.R.
<i>Philonicus albiceps</i> (Meigen)	L	England
<i>Satanas gigas</i> Eversmann	E, IL, P	Kazakhstan, U.S.S.R.
<i>Triorla interrupta</i> (Macquart) (as <i>Erax maculatus</i> Macquart)	L	U.S.A.
<i>Triorla interrupta</i> (Macquart) (as <i>Erax interruptus</i> Macquart)	L, P	Louisiana, U.S.A.
		Osterberger, 1930

Zinov'eva, 1959

Zinov'eva, 1959

Clements and Bennett, 1969

Copello, 1927, 1942

Cole and Pritchard, 1964

Cole and Pritchard, 1964

Peterson, 1957

Lavigne and Holland, 1969

Lehr, 1958

Davis, 1919

Peterson, 1957


Tsacas, 1969

Zinov'eva, 1959

Brindle, 1962


Zinov'eva, 1959

Davis, 1919



FIGS. 2-5. *Neomochtherus angustipennis* (Hine); pupal skin: 2, ventral view; 3, lateral view; 4, posterolateral view of last abdominal segment; 5, dorsal view. aap, anterior antennal process; acs, anterior coxal sheath; amsp, anterior mesothoracic spines; hs, hypopharyngeal sheath; ls, labral sheath; ms, maxillary sheath; pap, posterior antennal process; pmsp, posterior mesothoracic spine.

processes (aap) not joined at base and a group of 3 basally fused posterior antennal processes (pap) located ventrolaterally on either side, the 2 outermost processes fused basally for a greater distance and thus appearing shorter than the innermost process. Outermost process rounded apically, the 2 innermost processes acute. Labral sheath (ls) with a slight keel apically that is rugulose on either side. Hypopharyngeal sheath (hs) with a minute tubercle on either side. Maxillary (ms) and anterior coxal sheaths (acs) entirely smooth. Paired prothoracic spiracles elongate-oval, surrounded by a ring of thickened cuticle basally, situated midlaterally at anterior margin of thorax. A pair of anterior mesothoracic spines (amsp), 1 short and blunt, the other long and sharply curved,



Aushman '71

FIGS. 6-9. *Neomochtherus angustipennis* (Hine): 6, Wing; 7, Head; 8, Male terminalia, dorsal view; 9, Same, lateral view.

on either side of thorax above bases of sheaths of second pair of legs. A short, dull posterior mesothoracic spine (pmsp) on tubercle at base of each wing sheath. No rugulose area on thorax above wing sheaths. Sheaths of third pair of legs reaching slightly beyond middle of abdominal segment 3. First abdominal segment with a transverse row of 10 long, subequal, apically recurved spines dorsally along anterior margin and 4 yellowish bristles behind lateral spiracle, venter obscured. Second segment with a median transverse row of alternate long and short, straight spines dorsally, a short row of 4 or 5 bristles dorsolaterally, 4 to 7 bristles behind the lateral spiracle, and a mesally interrupted row of bristles ventrally. Third through seventh abdominal segments similar,

each with spines and bristles dorsally as on second segment, 5 or 6 bristles laterally, and complete ventral transverse row of bristles. Abdominal segment 8 (last segment) composed of a ringlike anterior portion with 4 subequal spines dorsally, 6 or 7 bristles laterally, and 12 bristles ventrally. Tapered posterior portion with a pair of long dorso-lateral processes, a pair of shorter ventrolateral processes, and a pair of short ventromedian processes. A pair of low tubercles midventrally on posterior part of segment 8.

The specimen is labeled, "Barcroft, Va., VIII.1.34, emerged VIII.13.34, pupa in surface soil under pine tree, J. C. Bridwell."

The adult compares exactly with Hine's (1909) description and figures, and with a male cotype from St. Elmo, Virginia, collected by F. C. Pratt, September 13 (USNM Type Number 12648). There is an additional adult in the Smithsonian Institution, a male, from Camp Peary, Virginia, July 1943, R. M. and G. E. Bohart. The Catalog of North American Diptera (Stone, et al., 1965) also records the species from North Carolina.

This species was described in the genus *Asilus* L. by Hine (1909), listed under *Asilus* in the North American Catalog, and under *Neomochtherus* by Hull (1962). Except for the rather well-developed facial gibbosity, *N. angustipennis* has the characters of the genus as discussed by Hull (1962) and Tsacas (1963, 1968). The only figures of this species that have been published are Hine's (1909) rather superficial drawings of the male terminalia; the head, wing, and male terminalia are shown here in Figures 6-9.

ACKNOWLEDGMENTS

Figure 1 was prepared by V. Krantz, Smithsonian Institution Photographic Services, and Figures 2-9 were prepared by A. D. Cushman, Systematic Entomology Laboratory, USDA.

I am very grateful to the following colleagues who read the manuscript and offered many suggestions for its improvement: A. N. Clements, Woodstock Agricultural Research Centre; R. J. Lavigne, University of Wyoming; H. Oldroyd, British Museum (N. H.); and L. Tsacas, Museum National d'Histoire Naturelle, Paris.

LITERATURE CITED

- ADAMOVIC, Z. R. 1963. Ecology of some asilid-species (Asilidae, Diptera) and their relation to honeybee (*Apis mellifera* L.). Mus. Hist. Nat. Beograd. 30: 1-104.
- . 1964. Feeding regime of some species of Asilidae (Diptera) in Yugoslavia. Arh. Biol. Nauka. 15(1-2): 37-74.
- ARTIGAS, J. N. 1970. Los Asilidos de Chile. Gayana, Zoologia 17: 1-473.
- BRAUNS, A. 1954a. Terricole Dipterenlarven. Musterschmidt, Göttingen. 179 pp.

- . 1954b. Puppen terricole Dipterenlarven. Musterschmidt, Göttingen. 156 pp.
- BRINDLE, A. 1962. Taxonomic notes on the larvae of British Diptera. 10. The Asilidae. Entomologist 95: 241-247.
- . 1968. Taxonomic notes on the larvae of British Diptera. 25. The larva and pupa of *Dioctria rufipes* (DeGeer) (Asilidae). Entomologist 101: 213-216.
- . 1969. Taxonomic notes on the larvae of British Diptera. 26. The presumed larva of *Dioctria oelandica* (Linnaeus) (Asilidae). Entomologist 102: 3-6.
- BROMLEY, S. W. 1930. Bee-killing robber flies. Jour. N. Y. Entomol. Soc. 38: 159-176.
- CARRERA, M., AND M. A. VULCANO. 1961. Relacao de alguns Asilidae (Diptera) e suas presas (4). Rev. Brasil de Entomol. 10: 67-80.
- CHAMPLAIN, A. B., AND J. N. KNULL. 1923. Notes on Pennsylvania Diptera. Entomol. News 34: 211-215.
- CLEMENTS, A. N., AND F. D. BENNETT. 1969. The structure and biology of a new species of *Mallophora* Macq. (Diptera, Asilidae) from Trinidad, W. I. Bull. Entomol. Res. 58: 455-463.
- COLE, F. R., AND A. E. PRITCHARD. 1964. The genus *Mallophora* and related asilid genera in North America (Diptera: Asilidae). Univ. Calif. Publ. Entomol. 36(2): 43-100.
- COPELLO, A. 1927. Biología del Moscardón cazador de abejas (*Mallophóra ruficauda*, Wied.). Argentina Min. Agri., Sec. Propag. e Inform., Circ. no. 699, 19 pp.
- . 1942. Moscardón cazador de abejas "*Mallophóra Ruficauda*." Rev. Apicultura, Argentina 19: no. 218, 4-5; 219, 10-11; 200, 13-14; 221, 14; 222, 12-13.
- CROUZEL, I. S. DE. 1965. Parasitismo en "Gusanos blancos" en la Republica Argentina (Col. Scarabeidae). Rev. Soc. Entomol. Argentina 27: 83-87.
- CUTHBERTSON, A. 1937. Biological notes on some Diptera in Southern Rhodesia. Proc. and Trans. Rhodesia Sci. Assn. 35: 16-34.
- DANIELS, N. E. 1966. The association of a rangeland grub, *Phyllophaga koehleriana* (Coleoptera: Scarabaeidae), with asilid larvae and with mites. Ann. Entomol. Soc. Amer. 59: 1021.
- DAVIS, J. J. 1919. Contributions to a knowledge of the natural enemies of *Phyllophaga*. Bull. Ill. Nat. Hist. Surv. 13: 53-138.
- DE GAVOTTO, A. L. R. 1964. Ciclo biológico de *Cyclocephala signaticollis* Burm. (Col., Scarabaeidae) y caracteres específicos de su larva. Rev. Invest. Agropec., ser. 5, Patol. Veg. 1: 151-161.
- ECIPENKO, P. A. 1967. The biology of *Eutolmus rufibarbis* Mg. (Diptera, Asilidae) in the Khabarovsk Region. Entomol. Obozr. 46: 109-112.

- FATTIG, P. W. 1944. The *Phyllophaga* or may beetles of Georgia. Emory Univ. Mus. Bull. 2: 1-32.
- . 1945. The Asilidae or robber flies of Georgia. Emory Univ. Mus. Bull. 3: 1-33.
- FELT, E. P. 1918. Grass and clover insects. Notes for the year. In 32nd report of the State Entomologist on injurious and other insects of the State of New York, 1916. N.Y. State Mus. Bull. No. 198, 276 pp.
- GREATHEAD, D. J. 1963. A review of the insect enemies of Acridoidea (Orthoptera). Trans. Roy. Entomol. Soc. London 114: 437-517.
- HENNIG, W. 1952. Die Larvenformen der Dipteren. Pt. 3 Akademie-Verlag, Berlin. 628 pp.
- HINE, J. S. 1909. Robberflies of the genus *Asilus*. Ann. Entomol. Soc. Amer. 2: 136-172.
- HOBBY, B. M. 1930. The British species of Asilidae (Diptera) and their prey. Trans. Entomol. Soc. South of England 6: 1-42.
- . 1933. Supplementary list of the prey of Asilidae (Dipt.) Entomol. Soc. South London 1(3): 69-74.
- HULL, F. M. 1962. Robber Flies of the world. The genera of the family Asilidae. U.S. Nat. Mus. Bull. 224, pt. 1. Vol. 1, p. i-x, 1-430; vol. 2, p. 433-907.
- IRWIN-SMITH, V. 1923. Studies in life-histories of Australian Diptera Brachycera. Pt. II. Asilidae. No. 1. Catalogue of the species of Asilidae of which the earlier stages have been recorded, with literary index, and bibliography of biological literature. Proc. Linn. Soc. N. S. Wales 48: 368-374.
- IWATA, K., AND A. NAGATOMI. 1962. Prey of some Japanese Asilidae. Kontyû 30: 87-100.
- KINOSHITA, E. 1940. Untersuchungen über die yesoraubfliege *Promachus yesonicus* Bigot unter berücksichtigung des forstschutzes. Hokkaido Imp. Univ., Fac. Agr., Expt. Forests Res. Bull. extra no. 1940: 171-270.
- LAVIGNE, R. J. 1963. Description of the eggs of *Leptogaster salvia* Martin (Diptera: Asilidae). Pan-Pac. Entomol. 39: 176.
- . 1964. Notes on the distribution and ethology of *Efferia bicaudata* (Diptera: Asilidae) with a description of the eggs. Ann. Entomol. Soc. Amer. 57: 341-344.
- , AND F. R. HOLLAND. 1969. Comparative behavior of eleven species of Wyoming robber flies (Diptera: Asilidae). Univ. Wyoming Agric. Exp. Sta. Sci. Monog. 18: 1-62.
- LEHR, P. A. 1958. The biology of *Promachus leontochlaenus* Loew (Diptera - Asilidae). Kazakh. Akad. Sel'sk. Nauk. Nauch.-Issled. Inst. Zash. Rast., Trudy 4: 210-224.
- . 1962. On the systematics and biology of *Eutolmus implacidus*

- Loew (Diptera, Asilidae). Akad. Nauk Kazakhskoi SSR Inst. Zool. Trudy 18: 205-215.
- . 1969. Ecologo-morphological analysis of robber flies (Diptera, Asilidae), 2. Entomol. Obozr. 48: 532-560.
- LINSLEY, E. G. 1960. Ethology of some bee- and wasp-killing robber flies of southeastern Arizona and western New Mexico (Diptera: Asilidae). Univ. Calif. Pub. Entomol. 16(7): 357-381.
- MELIN, D. 1923. Contributions to the knowledge of the biology, metamorphosis and distribution of the Swedish asilids in relation to the whole family of asilids. Zool. Bidrag Uppsala 8: 1-317.
- MUSO, J. J. 1967. Etude de la nymphe d'*Andrenosoma bayardi* Séguin, 1952 (Dipt., Asilidae). Bull. Soc. Entomol. France 72: 75-80.
- OSTERBERGER, B. A. 1930. *Erax interruptus* Macq. as a predator. Journ. Econ. Entomol. 23: 709-711.
- PARMENTER, L. 1952. The pupa of *Isopogon brevis* Meigen (Asilidae). Proc. Roy. Entomol. Soc. London, ser. A, 27: 43-44.
- PETERSON, A. 1957. Larvae of insects. Part II. Coleoptera, Diptera, Neuroptera, Siphonaptera, Mecoptera, Trichoptera. Edwards Brothers, Ann Arbor, Michigan. v + 416 pp.
- POULTON, E. B. 1924. The relation between the larvae of the asilid genus *Hyperechia* and those of xylocopid bees. Trans. Entomol. Soc. London 1924: 121-133.
- POWELL, J. A., AND G. I. STAGE. 1962. Prey selection by robber flies of the genus *Stenopogon*, with particular observations on *S. engelhardti* Bromley (Diptera: Asilidae). Wasmann Jour. Biol. 20: 139-157.
- QUENTIN, R. M. 1948. Sur la nymphe de *Laphria gilva* L. (Dipt. Asilidae). L'Entomologiste 4: 9-10.
- REINHARD, E. G. 1938. The egg-laying and early stages of the robber fly, *Erax aestuans* (Diptera: Asilidae). Entomol. News 49: 281-283.
- RITCHER, P. O. 1940. Kentucky white grubs. Ky. Agr. Expt. Sta. Bull. 401: 73-157 (Ky. Agr. Expt. Sta. Ann. Rpt. 53: 73-157).
- SCHMID, J. M. 1969. *Laphria gilva* (Diptera: Asilidae), a predator of *Dendroctonus ponderosae* in the Black Hills of South Dakota. Ann. Entomol. Soc. Amer. 62: 1237-1241.
- STONE, A., ET AL. 1965. A catalog of the Diptera of America north of Mexico. U.S.D.A. Agr. Hdbk. No. 276. p. i-iv, 1-1696. U.S. Gov. Print. Off., Washington.
- STOWER, W. J., G. B. POPOV, AND D. J. GREATHEAD. 1958. Oviposition behaviour and egg mortality of the Desert Locust (*Schistocerca gregaria* Forskål) on the coast of Eritrea. Anti-Locust Bull. 30: 1-33.
- STUBBS, A. E. 1970. Observations on the oviposition of *Leptarthus*

- brevirostris* (Mg.) (Diptera: Asilidae). *The Entomol.* 103: 289-293.
- TSACAS, L. 1963. Révision des *Neomochtherus* Osten-Sacken (Diptera: Asilidae) décrits par Macquart. *Bull. Mus. Natl. d'Hist. Nat.* 35: 257-272.
- . 1968. Révision des espèces du genre *Neomochtherus* Osten-Sacken. Diptères: Asilidae. I. Région Paléarctique. *Mém. Mus. Natl. Hist. Natl. N. S. ser. A. Zool.* 47(3): 128-328, pl. I-VIII, carte I-XI.
- . 1969. Révision des espèces du genre *Neomochtherus* Osten-Sacken (Diptères: Asilidae) II. Région éthiopienne. *Mém. Mus. Natl. d'Hist. Natl. N. S. ser. A, Zool.* 61(1): 1-32.
- , R. DESMIER DE CHENON, AND R. COUTIN. 1970. Observations sur le parasitisme larvaire d'*Hyperechia bomboidea* (Dipt. Asilidae). *Ann. Soc. Entomol. France (N. S.)* 6(2): 493-512.
- WEINBERG, M. 1968. Sur la biologie de *Dysmachus fuscipennis* (Mg.) (Diptera, Asilidae). *Trav. Mus. d'Hist. Nat. "Gr. Antipa"* 8(2): 875-884.
- WICHMANN, H. E. 1956. Untersuchungen über *Ips typographus* L. und seine Umwelt. Asilidae, Raubfliegen. *Zeitschr. F. Angew. Entomol.* 39: 58-62.
- ZINOV'eva, L. A. 1959. The biology and ecology of robber flies (Diptera, Asilidae) in the Naryn Sands (West Kazakhstan). *Entomol. Obozr.* 38: 554-567.